



A WNS COMPANY

2025 Computed Tomography (CT) Cervical Spine

Diagnostic Imaging

CT-CSpine-HH

Copyright © 2025 WNS (Holdings) Ltd.

Last Review Date: 03/24/2025

Previous Review Date: 10/28/2024

Guideline Initiated: 06/30/2019



Table of Contents

Computed Tomography (CT) Cervical Spine	3
CT General Contraindications	3
Preamble: Pediatric Diagnostic Imaging	3
CT Cervical Spine Guideline	3
CT Myelogram Guideline	7
Combination CT and MRI for Metastases Evaluation Guideline	8
Combination CT Cervical Spine and MRI Cervical Spine Guideline	8
Blood/Bone Marrow Cancers Surveillance section	9
Acute Lymphoblastic Leukemia Surveillance	9
Acute Myeloid Leukemia Surveillance reuse	9
Chronic Lymphocytic Leukemia/Small Cell Lymphocytic Lymphoma Surveillance ...	9
Chronic Myeloid Leukemia Surveillance	9
Hairy Cell Leukemia Surveillance	9
Multiple Myeloma Surveillance	9
Spine Surveillance section	10
Bone Cancer Surveillance	10
Central Nervous System (CNS) Cancer Surveillance	11
Neuroendocrine and Adrenal Tumors Surveillance	12
Occult Primary Cancer Surveillance	17
CT Cervical Spine Summary of Changes	17
CT Cervical Spine Procedure Codes	17
CT Cervical Spine Definitions	17
CT Cervical Spine References	23
Disclaimer section	24
Purpose	24
Clinician Review	24
Payment	25
Registered Trademarks (®/™) and Copyright (©)	25
National and Local Coverage Determination (NCD and LCD)	25
Background	25
Medical Necessity Codes	26

Computed Tomography (CT) Cervical Spine

**NCD 220.1**

See also, **NCD 220.1**: Computed Tomography at <https://www.cms.gov/medicare-coverage-database/search.aspx> if applicable to individual's healthplan membership.

CT General Contraindications

Computed tomography (CT) is contraindicated (relative) for **ANY** of the following:

1. Allergy/idiosyncratic reaction to contrast material (if intravascular contrast material is used)
2. Pregnancy
3. Renal impairment (glomerular filtration rate [GFR] is less than 30 ml/min/1.73 m².)

References: [3]

Preamble: Pediatric Diagnostic Imaging

HealthHelp's clinical guidelines for the Diagnostic Imaging program, are intended to apply to both adults and pediatrics (21 years of age or younger), unless otherwise specified within the criteria.

CT Cervical Spine Guideline

Computed tomography (CT) of the cervical spine is considered medically appropriate when the documentation demonstrates **ANY** of the following: (***NOTE:** *When requested in combination with magnetic resonance imaging [MRI] cervical spine, need for soft tissue and bony anatomy evaluation must be required.*)

1. Cervical radiculopathy is demonstrated on electromyography (EMG) or nerve conduction study. (***NOTE:** *An EMG is **NOT** recommended to determine the cause of axial lumbar, cervical or thoracic spine pain.*)

References: [15]

2. Pain in the cervical neck area, MRI is **contraindicated or unavailable** and **ANY** of the following: [15]
 - a. Conservative management (chiropractic treatments, physical therapy) and **EITHER** of the following:
 - i. Attempted within the last 6 months, for at least 6 weeks **AND** symptoms persist or worsen.

- ii. Symptoms progress or worsen during course of conservative management
- b. Neck pain is isolated, in a pediatric individual and **ANY** of the following red flags:
(***NOTE**: *Conservative care is not required if red flags are present.*)
 - i. Age is 5 years or younger.
 - ii. Fever, malaise or weight loss (5% in 1 month or 10% in 6 months)
 - iii. Neurologic examination is abnormal.
 - iv. Pain at night disrupts or prevents sleep.
 - v. Pain is constant.
 - vi. Pain lasts more than 4 weeks.
 - vii. Postural changes (eg, kyphosis or scoliosis)
 - viii. Radicular pain
 - ix. Stiffness/gelling in early morning

References: [7]

- 3. Arnold-Chiari malformation is known and MRI is **contraindicated or unavailable**.

References: [18]

- 4. Cancer, tumor or metastasis evaluation, MRI is **contraindicated or unavailable** and **ANY** of the following: (***NOTE**: *CT is also needed to further characterize solitary indeterminate lesions seen on MRI.*)
 - a. Prior imaging for metastasis or tumor is abnormal, non-diagnostic or indeterminate.
 - b. Spinal tumor is known **AND** signs are new or progressing (eg, non-traumatic pain is new or increasing).
 - c. Surveillance following the **National Comprehensive Cancer Network (NCCN) Guideline's** surveillance recommendations (see **Surveillance** section).

References: [6] [14]

- 5. Cerebrospinal fluid (CSF) leak is suspected (eg, cerebrospinal-venous fistula, orthostatic headache, otorrhea, post lumbar puncture headache, post spinal surgery headache, rhinorrhea, spontaneous idiopathic intracranial hypotension [SIH]) **AND** MRI is **contraindicated or unavailable**. (***NOTE**: *for cerebrospinal-venous fistula, CT Myelogram is preferred.*)

References: [9]

- 6. Compression fracture(s) evaluation and **ANY** of the following:

- a. **ALL** of the following:
 - i. Demonstrated on X-ray
 - ii. Neck pain is worsening.
 - iii. New
 - iv. **NO** known malignancy
- b. Compression fractures are known and treated **AND** neck pain is new.

References: [14]

7. Infection (eg, abscess, discitis, osteomyelitis) is suspected or known, MRI is **contraindicated or unavailable** and **ANY** of the following:
 - a. Active treatment, to assess response
 - b. Immune system suppression-related (eg, cancer, diabetes, dialysis, human immunodeficiency virus [HIV], intravenous drug use) spinal infection is suspected, from signs (eg, abnormal white blood cell count, erythrocyte sedimentation rate [ESR], back pain).
 - c. Signs/symptoms are present (eg, chills, complete blood count [CBC], c-reactive protein [CRP], ESR, fever, pain)

References: [17] [15]

8. Inflammatory disease or atlantoaxial instability is suspected or known **AND** MRI is **contraindicated or unavailable** **OR** for surgical treatment planning and **ANY** of the following:
 - a. Atlantoaxial articulation disorder (eg, Down syndrome, Marfan syndrome) is known and **ANY** of the following:
 - i. Cervical spine X-ray(s) are abnormal, non-diagnostic or indeterminate.
 - ii. Neurological physical exam is abnormal (eg, abnormal gait or reflexes, bowel/bladder dysfunction, extremity weakness).
 - b. Cervical instability is suspected, based on symptoms (eg, dizziness, numbness, pain, tingling) **AND** X-ray is negative.
 - c. Neuroinflammatory conditions (Behcet's syndrome, sarcoidosis) are suspected **AND** neurological exam is abnormal and rheumatology evaluation (eg, CRP, ESR) is completed.
 - d. Rheumatoid arthritis with neurologic symptoms (eg, dizziness, numbness, pain, tingling) **OR** X-ray demonstrates subluxation. (***NOTE:** Initial imaging should be a lateral X-ray in flexion and neutral.)

- e. Spondyloarthropathies are suspected or known, when X-ray(s) are non-diagnostic or indeterminate and rheumatology evaluation (eg, CRP, ESR) is completed.

References: [8] [12] [4]

- 9. Myelopathy is suspected, MRI is **contraindicated or unavailable** and **ANY** of the following: (***NOTE:** *Conservative care is **NOT** required prior to ordering imaging.*)
 - a. Abnormal neurologic physical exam (eg, abnormal gait or reflexes, bowel/bladder dysfunction, extremity weakness) is demonstrated on physical exam (eg, digital rectal exam, examination of balance and reflexes)
 - b. Symptoms or neurologic physical exam (eg, balance, difficulty with ambulation, diffuse numbness in the hands, grasping and holding objects, hand clumsiness, pins and needles sensation) are progressing.

References: [1]

- 10. Neurological deficits/symptoms (eg, abnormal reflexes, loss of sensation, numbness/tingling) are known based on completed neurological exams (eg, Lhermitte's sign, shoulder abduction test, spurling test) and MRI is **contraindicated or unavailable**.

References: [20]

- 11. Post-surgical assessments for evaluation of complications or disease recurrence.
- 12. Scoliosis evaluation, MRI is **contraindicated or unavailable** and **ANY** of the following:
 - a. Atypical curve (eg, Kyphosis more than 30 degrees, left thoracic curve, short segment)
 - b. Congenital scoliosis or juvenile idiopathic scoliosis and age is less than 10, for initial assessment.
 - c. Neurological deficit is new or unexplained.
 - d. Pre-operative planning
 - e. Spinal deformity is progressive.
 - f. Treatment planning depends on imaging

References: [11]

- 13. Syring or syringomyelia is suspected or known, MRI is **contraindicated or unavailable** and **ANY** of the following:
 - a. Predisposing conditions are known (eg, Arnold-Chiari malformation, neoplasm, prior trauma, severe spondylosis).
 - b. Prior imaging demonstrates an abnormality. (eg, deformities, nodules, septations)

14. Tethered cord or spinal dysraphism is suspected or known, from preliminary imaging, neurological exam and/or high cutaneous stigmata **AND** MRI is **contraindicated or unavailable**.
15. Toe walking, in a pediatric individual, with signs/symptoms of myelopathy (eg, numbness, pain, tingling) localized to the cervical spine **AND** MRI is **contraindicated or unavailable**.
16. Trauma or acute injury evaluation with **ANY** of the following:
 - a. National emergency X-Radiography utilization study (NEXUS) or Canadian cervical rules (CCR) criteria for imaging are met for **ANY** of the following: (***NOTE:** *CT and MRI provide complementary information. When indicated it is appropriate to perform both.*)
 - i. Evaluation with CT for initial imaging
 - ii. Spine is unstable, CT or MRI for treatment planning.
 - iii. Spinal cord or nerve root injury is suspected **OR** when obtunded and CT is negative.
 - b. Nerve root injury is suspected.
 - c. Spinal abnormalities (eg, ankylosing spondylitis, diffuse idiopathic skeletal hyperostosis) are known. (***NOTE:** *Both a MRI and CT are appropriate.*)
 - d. Unexaminable condition (eg, distracting injury, Glasgow coma scale is less than 15)

References: [5] [13] [16] [19]

CT Myelogram Guideline

Computed tomography (CT) myelogram is considered medically appropriate when the documentation demonstrates a **MRI is contraindicated or unavailable OR** it is the surgeon's preference, and **ANY** of the following:

1. Brachial plexus (if appropriate) **OR** nerve root injury is suspected in the neonate.
2. Cerebral spinal fluid (CSF) leak is known (eg, from symptoms of headache after lumbar puncture, post spinal surgery, spontaneous intracranial hypertension [SIH]), to demonstrate location of leak.
3. Magnetic resonance imaging (MRI) findings are incongruent with symptoms.
4. Pre-procedural planning (eg, nerve roots or dural sac evaluation)

References: [10] [2019 ACR-ASNR-SPR Practice Parameter for the Performance of Myelography and Cisternography]

Combination CT and MRI for Metastases Evaluation Guideline

Combination CT/MRI studies (5 or less concurrent studies, with a CT or MRI appropriate for cancer location: abdomen, brain, cervical spine, chest, lumbar spine, neck, pelvis and/or thoracic spine) for **ANY** of the following situations:

1. Staging evaluation, for baseline pre-therapy
2. Surveillance following the National Comprehensive Cancer Network (NCCN) Guidelines recommended schedule (See **Surveillance** section)

Combination CT Cervical Spine and MRI Cervical Spine Guideline

Computed tomography (CT) cervical spine **combined** with magnetic resonance imaging (MRI) cervical spine is considered medically appropriate when the documentation demonstrates the following: (***NOTE: NOT** an all inclusive list)

1. Bony and soft tissue abnormality is known (eg, deformity, prior imaging, physical exam) **AND** imaging may change the treatment plan.
2. Craniocervical junction is unstable (based on prior imaging or pronounced bulbar or myelopathic signs).
3. Fractures are complex or pathological.
4. Malignant process of spine is suspected or known (eg, neurological deficits, worsening back pain), with **BOTH** bony and soft tissue involvement.
5. Ossification of posterior longitudinal ligament (OPLL) is suspected (eg, myelopathy, radiculopathy, spinal cord compression) **OR** for re-evaluation of known OPLL.



LCD 35175

See also, **LCD 35175**: MRI and CT Scans of the Head and Neck at <https://www.cms.gov/medicare-coverage-database/search.aspx> if applicable to individual's healthplan membership.



LCD 35391

See also, **LCD 35391**: Multiple Imaging in Oncology at <https://www.cms.gov/medicare-coverage-database/search.aspx> if applicable to individual's healthplan membership.

**LCD 37373**

See also, **LCD 37373**: MRI and CT Scans of Head and Neck at <https://www.cms.gov/medicare-coverage-database/search.aspx> if applicable to individual's healthplan membership.

Blood/Bone Marrow Cancers Surveillance section

Acute Lymphoblastic Leukemia Surveillance

Acute lymphoblastic leukemia: No imaging surveillance suggested.

References: [2024 Acute Lymphoblastic Leukemia Version 3.2024]

Acute Myeloid Leukemia Surveillance reuse

Blastic plasmacytoid dendritic cell neoplasm surveillance includes a repeat PET/CT for individuals with prior evidence of extramedullary disease.

References: [2025 Acute Myeloid Leukemia (Age ≥ 18) Version 2.2025]

Chronic Lymphocytic Leukemia/Small Cell Lymphocytic Lymphoma Surveillance

Chronic lymphocytic leukemia/small cell lymphocytic lymphoma: No imaging surveillance suggested.

References: [2025 Chronic Lymphocytic Leukemia/Small Lymphocytic Leukemia Version 1.2025]

Chronic Myeloid Leukemia Surveillance

Chronic Myeloid Leukemia: No imaging surveillance suggested.

References: [2025 Chronic Myeloid Leukemia Version 3.2024]

Hairy Cell Leukemia Surveillance

Hairy cell leukemia: No imaging surveillance suggested.

References: [2025 Hairy Cell Leukemia Version 1.2025]

Multiple Myeloma Surveillance

Multiple myeloma surveillance includes **ANY** of the following:

1. Multiple myeloma, surveillance imaging when recurrence is suspected with **ANY** of the following:

- a. CT scan, low dose
 - b. FDG PET/CT
 - c. MRI (- contrast material), whole-body
2. Smoldering myeloma, surveillance imaging annually (or more often when recurrence is suspected) with **ANY** of the following:
 - a. CT scan, low dose
 - b. FDG PET/CT
 - c. MRI (- contrast material), whole-body

References: [2025 Multiple Myeloma Version 2.2025]

Spine Surveillance section

Bone Cancer Surveillance

Bone cancer surveillance includes **ANY** of the following:

1. Chondrosarcoma surveillance for **ANY** of the following:
 - a. Atypical cartilaginous tumor surveillance with cross-sectional imaging (CT + contrast, MRI \pm contrast) every 6 to 12 months for 2 years, then annually as clinically indicated
 - b. Low-grade, extracompartmental appendicular tumor, grade I axial tumors or high-grade (grade II or III, clear cell or extracompartmental) tumors surveillance with **ALL** of the following:
 - i. Chest CT at least every 6 months for 5 years, then annually for at least 10 years, then if symptoms are new or progressing.
 - ii. MRI (\pm contrast) or CT (+ contrast) if symptoms are new or progressing.
2. Chordoma surveillance with **ALL** of the following:
 - a. Chest CT imaging every 6 months, annually for 5 years, then annually thereafter, then if symptoms are new or worsening.
 - b. Imaging of primary site, timing and modality (eg, MRI \pm CT [both + contrast]) if symptoms are new or progressing, up to 10 years
3. Ewing Sarcoma after primary treatment completed surveillance with **ALL** of the following:
 - a. Chest CT: every 3 months

- b. Primary site imaging with MRI ± CT (both + contrast), increase intervals after 24 months and after 5 years, annually, then if symptoms are new or progressing (indefinitely) (***NOTE: PET/CT [head-to-toe] is appropriate**)
4. Giant cell tumor of the bone surveillance with **ALL** of the following:
 - a. Chest CT or MRI imaging every 6 to 12 months for 4 years, then annually thereafter, then if symptoms are new or progressing
 - b. Surgical site imaging if symptoms are new or progressing (eg, CT and/or MRI, both with contrast)
5. Osteosarcoma surveillance with primary site and chest imaging (using same imaging that was done for initial work-up) for **ANY** of the following: (***NOTE: PET/CT [head-to-toe] is appropriate.**)
 - a. Image every 3 months for years 1 and 2
 - b. Image every 4 months for year 3
 - c. Image every 6 months for years 4 and 5
 - d. Image annually for year 6 and thereafter, then if symptoms are new or progressing

References: [2025 Bone Cancer Version 2.2025]

Central Nervous System (CNS) Cancer Surveillance

Central nervous system (CNS) cancer surveillance includes **ANY** of the following:

1. Glioblastoma, *IDH* wild-type, magnetic resonance imaging with (MRI) of the brain and **ANY** of the following:
 - a. Pre-operative and post-operative; within 48 hours
 - b. Pre-radiation planning; every 3 to 5 weeks, post-operatively
 - c. Post-radiation; 3 to 6 weeks post-radiation, then every 2 to 3 months for 3 years, then every 2 to 4 months indefinitely
2. Glioma, imaging with MRI of the brain and **ANY** of the following:
 - a. Astrocytoma, *IDH* mutated and **ANY** of the following:
 - i. Grade 2 and **ANY** of the following:
 - A. After radiation therapy (RT) **AND** chemotherapy: every 6 months until tumor progression
 - B. After RT **OR** chemotherapy: every 3 to 4 months for the 1st 5 years, then every 3 to 4 months until tumor progression

- C. After surgery: every 3 to 4 months until tumor progression
 - ii. Grade 3 and **ANY** of the following;
 - A. After RT **AND** chemotherapy: every 6 months until tumor progression
 - B. After RT **OR** chemotherapy: every 3 to 4 months for the 1st 5 years, then every 3 to 4 months until tumor progression
 - iii. Grade 2 or 3, recurrent; image every 2 to 3 months
- b. Oligodendroglioma, *IDH* mutated, 1p/19q co-deleted and **ANY** of the following:
 - i. Grade 2 and **ANY** of the following:
 - A. After radiation therapy (RT) **AND** chemotherapy: every 6 to 9 months until tumor progression
 - B. After RT **OR** chemotherapy: every 3 to 4 months for the 1st 5 years, then every 3 to 4 months until tumor progression
 - C. After surgery: every 3 to 4 months until tumor progression
(***NOTE:** For individuals who underwent gross total resection, every 6 to 9 months for 5 years post-surgery until tumor progression)
 - ii. Grade 3 and **ANY** of the following:
 - A. After radiation therapy (RT) **AND** chemotherapy: every 6 to 9 months until tumor progression
 - B. After RT **OR** chemotherapy: every 3 to 4 months for the 1st 5 years, then every 3 to 4 months until tumor progression
 - iii. Grade 2 or 3, recurrent, image every 3 to 4 months
- 3. Leptomeningeal metastases imaging with MRI of the brain and/or total spine every 2 to 3 months for the 1st 2 years, every 6 months until year 5, then annually indefinitely
- 4. Medulloblastoma, imaging with MRI of the brain every 2 to 3 months for 2 years
- 5. Primary CNS lymphoma, image every 2 to 3 months for 2 years

References: [2025 Central Nervous System Cancers Version 1.2025]

Neuroendocrine and Adrenal Tumors Surveillance

Neuroendocrine and adrenal cancer surveillance includes **ANY** of the following:¹

¹**NO** surveillance is indicated for appendiceal tumors 2 cm or smaller **WITHOUT** aggressive features (eg, high-grade cytologic atypia, infiltrative invasion lymphatic and hematogenous metastases).

1. Adrenal gland tumors surveillance imaging includes **ANY** of the following:
 - a. Localized disease: chest computed tomography (CT) (\pm contrast) and abdominal CT or magnetic resonance imaging (MRI) (+ contrast) every 3 to 12 months up to 5 years, then if symptoms are new or progressing.
 - b. Locoregional unresectable or metastatic disease; chest CT (\pm contrast) and CT or MRI abdomen and pelvis (+ contrast) or FDG positron emission tomography (PET)/CT every 3 to 12 months up to 5 years, then if symptoms are new or progressing.
2. Carcinoid syndrome surveillance imaging includes **BOTH** of the following:
 - a. Abdominal/pelvic multiphasic CT or MRI every 3 to 12 months and chest CT (\pm contrast) if symptoms are new or progressing.
 - b. Echocardiogram (ECHO) every 1 to 3 years or as clinically indicated **WITHOUT** known carcinoid heart disease (CHD) and at least annually for individuals with established CHD.
3. Gastrointestinal tract (well-differentiated grade 1/2), lung and thymus imaging and **ANY** of the following:
 - a. Lung nodules, multiple or tumorlets, image with chest CT (- contrast) every 12 to 24 months if symptoms are new or progressing.
 - b. Rectal tumor is 1 cm to 2 cm or less: image with rectal MRI at 6 and 12 months if symptoms are new or progressing.
4. Gastrointestinal (GI) tract (jejunum/ileum/colon, duodenum, rectum), lung and/or thymus neuroendocrine tumor (NET) surveillance includes imaging post-resection with **ANY** of the following:
 - a. Jejunum/ileum/colon, duodenum, rectum and thymus, surveillance imaging with abdominal \pm pelvic multiphasic CT or MRI according to **ONE** of the following levels of frequency²:
 - i. Within 3 months to 12 months post-operatively
 - ii. After 12 months, image every 12 to 24 months for 10 years
 - iii. After 10 years if symptoms are new or progressing.
 - b. Lung/thymus tumors surveillance chest CT (\pm contrast) for primary tumors, (as clinically indicated for primary GI tumors) according to **ONE** of the following levels of frequency:

²High-grade tumors are appropriate for more frequent monitoring.

- i. Within 12 weeks to 12 months post-operatively
 - ii. After 12 months, image every 12 to 24 months for 10 years
 - iii. After 10 years if symptoms are new or progressing.
- 5. Grade 3, well-differentiated neuroendocrine surveillance includes chest CT (\pm contrast) as clinically indicated for **ANY** of the following:
 - a. Locally advanced/metastatic disease with favorable biology (low Ki-67 [eg, less than 55%], positive somastatin receptor [SSTR] based PET imaging) includes abdominal/pelvic MRI (+ contrast) or abdominal/pelvic multiphasic CT for surveillance with **ANY** of the following:
 - i. Resectable disease surveillance every 3 to 6 months for 2 years, then every 6 to 12 months for up to 10 years **AND** chest CT if symptoms are new or progressing.
 - ii. Unresectable disease surveillance every 12 weeks to 24 weeks (depending on tumor biology) **AND** chest CT (\pm contrast), SSTR-PET/CT, SSTR-PET/MRI or FDG-PET/CT; if symptoms are new or progressing.
 - b. Locally advanced/metastatic disease with unfavorable biology (high Ki-67 [eg 55% or higher], rapid growth rate, FDG avid tumors, negative SSTR-based PET imaging), includes surveillance imaging, every 8 weeks to 12 weeks (depending on tumor biology) with **ALL** of the following:
 - i. Abdominal/pelvic MRI (+ contrast) or abdominal/pelvic multiphasic CT and FDG PET/CT as clinically indicated
 - ii. Chest CT (\pm contrast) if symptoms are new or progressing.
 - iii. FDG-PET/CT, if symptoms are new or progressing.
 - c. Locoregional disease (resectable) abdominal/pelvic MRI (+ contrast) or abdominal/pelvic multiphasic CT with frequency of **ONE** of the following:
 - i. Every 3 to 6 months for 2 years (depending on tumor biology, Ki-67) and chest CT as clinically indicated
 - ii. Every 6 months to 12 months for up to 10 years (depending on tumor biology, Ki-67) and chest CT as clinically indicated
 - d. Multiple endocrine neoplasia, type 1 (MEN1) screening surveillance for **ANY** of the following tumor types: (***NOTE:** *For prolonged surveillance, use imaging studies without radiation.*)
 - i. Lung/thymic NETs: chest CT or MRI (+ contrast) every 1 to 3 years

- ii. PanNET: abdominal/pelvic CT or MRI (+ contrast) every 1 to 3 years
 - iii. Parathyroid: if calcium rises, re-image with single-photon emission computed tomography (SPECT) scan (SPECT-CT preferred) or 4D-CT
 - iv. Pituitary: pituitary or sella MRI (+ contrast) of the pituitary every 3 to 5 years
- e. Poorly differentiated large or small cell carcinoma and/or mixed neuroendocrine/non-neuroendocrine neoplasm or unknown primary, imaging surveillance includes **ALL** of the following:
 - i. Locoregional unresectable or metastatic disease surveillance imaging includes **EITHER** chest CT (\pm contrast) with abdominal/pelvic MRI (+ contrast) **OR** chest/abdominal/pelvic multiphasic CT; every 6 weeks to 16 weeks
 - ii. Resectable surveillance imaging includes **EITHER** chest CT (\pm contrast) with abdominal/pelvic MRI (+ contrast) **OR** chest, abdomen and pelvis multiphasic CT; every 12 weeks for the 1st year, and every 6 months thereafter
- f. Post-operative from potentially curative surgery surveillance for at least 10 years (longer if high-risk)
- 6. Pancreatic neuroendocrine tumor surveillance imaging, post-resection, includes chest CT (\pm contrast) as clinically indicated and abdominal multiphasic CT or MRI with imaging frequency of **ONE** of the following³:
 - a. Within 3 to 12 months post-operatively
 - b. After 12 months, image every 6 to 12 months for 10 years
 - c. After 10 years if symptoms are new or progressing.
- 7. Pheochromocytoma/paranganglioma surveillance imaging and **ANY** of the following:
 - a. Locally unresectable disease or distant metastases, imaging every 12 weeks for 12 months, includes **ANY** of the following:
 - i. Chest, abdomen and pelvis CT with contrast
 - ii. Chest CT (\pm contrast) and abdominal/pelvic MRI (- contrast) (if risk for hypertensive episode)
 - iii. FDG-PET/CT for bone dominant disease

³High-grade tumors are appropriate for more frequent monitoring.

- iv. SSTR-PET/CT or SSTR-PET/MRI (if previous SSTR-positive or concern for disease progression) prior to radionuclide therapy
- b. Resectable disease, post-resection includes chest CT (\pm contrast) and abdominal/pelvic CT or MRI (+ contrast), if clinically indicated with imaging frequency of **ONE** of the following:
 - i. 12 weeks to 12 months after resection
 - ii. Every 6 to 12 months for the 1st 3 years
 - iii. Annually from year 4 to 10.
 - iv. More than 10 years, then as clinically indicated



TIP

NCCN recommends following the surveillance protocols from designated guidelines for the following hereditary endocrine neoplasia syndromes :

- Thyroid cancer guideline, use for: Multiple endocrine neoplasia, type 2 (MEN2) with genetic evaluation of inherited syndromes
- Kidney cancer, use for:
 - Hereditary paraganglioma/pheochromocytoma syndrome
 - Tuberous sclerosis complex (TSC1 and TSC2)
 - von Hippel Lindau syndrome (VHL)
- Genetic/Familial High-Risk Assessment: Breast, Ovarian, and Pancreatic, use for:
 - Neurofibromatosis type 1 (NF1)
 - Li-Fraumeni syndrome (TP53)
 - Lynch syndrome (MLH1, EPCAM/MSH2, MSH6, PMS2)
- Genetic/Familial High-Risk Assessment: Colorectal, use for:
 - Lynch syndrome (MLH1, EPCAM/MSH2, MSH6, PMS2)
 - Familial adenomatous polyposis (APC)

References: [2025 Neuroendocrine and Adrenal Tumors Version 2.2025]

Occult Primary Cancer Surveillance

Occult primary cancer surveillance imaging for long-term surveillance includes diagnostic tests based on symptomatology.

References: [2025 Occult Primary Version 2.2025]

CT Cervical Spine Summary of Changes

CT Cervical Spine guideline had the following version changes from 2024 to 2025:

- Added "Glomerular filtration rate" to "Renal impairment" under Contraindications
- Change in wording for "Peri-procedural" indication
- Citations updated per the evidence
- Removed the following as current evidence no longer supports the indication:
 - Combination studies as they are redundant
 - "Limp in a young child (age is 5 years or younger)" as it is redundant with "Age is 5 years or younger" under "Pain in the cervical neck area" under "Neck pain is isolated"
- 6/23/2025 Mid-cycle update
 - Established parameters for "Active treatment"
 - Added time frame for "Post-surgical assessment"
 - Added examples throughout
 - Added parameters for "Arnold-chiari malformation"

CT Cervical Spine Procedure Codes

Table 1. CT Cervical Spine Associated Procedure Codes

CODE	DESCRIPTION
72125	Computed tomography, cervical spine; without contrast material
72126	Computed tomography, cervical spine; with contrast material
72127	Computed tomography, cervical spine; without contrast material, followed by contrast material(s) and further sections

CT Cervical Spine Definitions

Abscess is a swollen area within body tissue, containing an accumulation of pus.

Achondroplasia is a rare autosomal dominant condition characterized by impaired endochondral ossification, resulting in disproportionate short stature and abnormal growth of long bones, vertebrae, and several bones in the skull.

Ankylosing spondylosis (spondylitis) is a chronic inflammatory disease that affects the spine, sacroiliac joints and often other joints (such as the shoulder), and is marked by pain and stiffness.

Arachnoiditis is defined as painful inflammation of the arachnoid mater, one of the meninges, and the spinal nerves, often leading to chronic pain and neurological deficits.

Articulation is a joint or juncture between bones or cartilages in the skeleton of a vertebrate.

Atlantoaxial refers to the anatomical and functional relationship between the first cervical vertebra (atlas, C1) and the second cervical vertebra (axis, C2), primarily responsible for the rotation of the neck.

Babinski reflex, also known as the Babinski reflex, is a neuro-pathological response in the foot that occurs when the sole of the foot is firmly stroked and the big toe moves upward or toward the top of the foot. The other toes may also fan out.

Behcet's disease is a chronic, relapsing systemic vasculitis characterized by recurrent oral and genital ulcers, uveitis, and various other systemic manifestations.

Bone scan is a nuclear imaging procedure that examines the bones in the skeleton. It can help diagnose and track bone diseases, and can also be used to monitor the progress of certain treatments.

Canadian Cervical Rules (CCR) are a set of guidelines that help clinicians decide if cervical spine imaging is needed for trauma patients in the emergency department. The CCR are used to rule out cervical spine injuries in low-risk patients. An example can be found at: https://www.physio-pedia.com/Canadian_C-Spine_Rule

Cerebrospinal fluid (CSF) is a colorless liquid that is comparable to serum, is secreted from the blood into the lateral ventricles of the brain, and serves chiefly to maintain uniform pressure within the brain and spinal cord.

Cerebrospinal fluid (CSF) leak is a leak of cerebrospinal fluid that results from a hole or tear in the dura (the outermost layer of the meninges).

Chaddock reflex is a diagnostic reflex similar to the Babinski reflex. Chaddock's sign is present when stroking of the lateral malleolus causes extension of the great toe, indicating damage to the corticospinal tract.

Chiari malformation (Arnold-Chiari syndrome) is a congenital abnormality in which the lower surface of the cerebellum and the lower brain stem protrude into the spinal canal through the foramen magnum.

Compression fracture is a break in the vertebrae and can cause the vertebrae to collapse, making them shorter.

Computed tomography (CT) is an imaging test that uses X-rays to computer analysis to generate cross sectional images of the internal structures of the body that can be displayed in multiple planes.

Computed tomography (CT) myelogram is a diagnostic imaging test generally done by a radiologist. It uses a contrast dye and X-rays or computed tomography (CT) to look for problems in the spinal canal.

Compression is reducing in size, quantity or volume, as if by squeezing.

Congenital is a condition or trait present from birth.

Conservative management is an approach to treating pain utilizing non-surgical treatment options such as physical therapy, medication and injections, for a designated time, usually 4 to 6 weeks.

C-reactive protein (CRP) is a pentameric protein synthesized by the liver, whose level rises in response to inflammation.

Delayed union fracture a fracture that has not healed within the expected timeframe for a specific fracture location, typically taking longer than 3 to 4 months to show adequate radiographic signs of healing.

Diffuse idiopathic skeletal hyperostosis (DISH) is a condition that causes ligaments to become calcified and hard. It usually affects the ligament around the spine, but it can also affect other areas of the body where ligaments join to bone.

Diplegia is a type of paralysis that affects similar body parts on both sides of the body, such as both arms or both legs. It's the most common cause of paralysis in children, but can affect people of any age. Unlike other forms of paralysis, diplegia is unpredictable and may improve, worsen or change over time.

Discitis is an uncommon primary infection of the vertebral disc, specifically the nucleus pulposus, often involving the cartilaginous end plate and vertebral body, and is most commonly caused by *Staphylococcus aureus*.

Discogram is a radiograph of an intervertebral disk made after injection of a radiopaque substance.

Down syndrome is a congenital condition characterized especially by developmental delays, usually mild to moderate impairment in cognitive functioning, short stature, upward slanting eyes, a flattened nasal bridge, broad hands with short fingers and decreased muscle tone caused by trisomy of the human chromosome numbered 21.

Drop metastases are intradural extramedullary spinal metastases that arises from intracranial lesions.

Dysraphism refers to congenital malformations of the spine and spinal cord resulting from incomplete midline closure during embryologic development.

Electromyogram (EMG) is a diagnostic test that measures the electrical activity of muscles at rest and during contraction using a needle electrode inserted into the muscle.

Erythrocyte sedimentation rate (ESR) is a blood test that measures the rate at which red blood cells settle at the bottom of a test tube over one hour, indicating the presence of inflammation in the body.

Fistula is an abnormal connection between two epithelialized surfaces, often involving organs such as the gut, bladder, vagina, or skin, and can result from various causes including surgery, trauma, Crohn's disease, diverticular disease, or malignancy.

Glasgow Coma Scale (GCS) is a clinical tool used to assess a patient's level of consciousness, particularly after traumatic brain injury (TBI). The GCS evaluates three aspects of responsiveness: eye opening, verbal response, and motor response. Scores range from 3 to 15, with higher scores indicating better neurological function.

Hemiparesis is muscular weakness or partial paralysis restricted to one side of the body.

Hoffmann's sign, also known as Hoffmann's reflex, is a neurological exam that involves flicking a patient's middle fingernail to see if their thumb or index finger flexes involuntarily. A positive result, also known as hyperreflexia, indicates that the nervous system is overreacting to the flick and upper motor neuron lesion or corticospinal pathway dysfunction may be present.

Human Immunodeficiency Virus (HIV) is a retrovirus that primarily infects CD4+ T lymphocytes, leading to progressive immunodeficiency and potentially resulting in AIDS.

Immunosuppression is the deliberate reduction or inhibition of the immune system's ability to respond to antigens, typically achieved through medications or therapies, to prevent organ rejection or treat autoimmune diseases, but it increases the risk of infections and malignancies.

Indeterminate findings are inconclusive or insufficient for treatment planning.

Kyphosis is the exaggerated outward curvature of the thoracic region of the spine resulting in a rounded upper back.

Leukemia is a type of cancer that affects the blood and bone marrow. It's caused by the rapid production of abnormal white blood cells. These abnormal white blood cells can't fight infection and make it harder for the bone marrow to produce red blood cells and platelets.

Leptomeningeal carcinomatosis is a severe complication of late-stage cancer characterized by the spread of malignant cells to the leptomeninges, including the pia mater, arachnoid, and subarachnoid space, leading to rapid mortality despite treatment.

Lhermitte's sign is an electric-like pain or tingling down the spine or limbs triggered by neck flexion or extension.

Lumbar puncture is a puncture of the subarachnoid space in the lumbar region of the spinal cord to withdraw cerebrospinal fluid or inject medications.

Lymphoma is a type of blood cancer that affects the immune system. Lymphoma occurs when abnormal white blood cells, called lymphocytes, grow in the lymphatic system.

Magnetic resonance imaging (MRI) is a non-invasive diagnostic technique that produces computerized images of internal body tissues and is based on nuclear magnetic resonance of atoms within the body induced by the application of radio waves.

Malaise is an indefinite feeling of debility or lack of health often indicative of or accompanying the onset of an illness.

Marfan syndrome is a disorder of connective tissue inherited as a dominant trait, characterized by abnormal elongation of the long bones and often with ocular and circulatory defects.

Metastases is the spread of a disease-producing agency (such as cancer cells) from the initial or primary site of disease to another part of the body.

Myelogram is a radiographic visualization of the spinal cord after injection of a contrast medium into the spinal subarachnoid space.

Myelopathy is a disease or disorder of the spinal cord or bone marrow.

National Emergency X-ray Utilization Study (NEXUS) is a very large, federally supported, multi-center, prospective study designed to define the sensitivity, for detecting significant cervical spine injury, of criteria previously shown to have high negative predictive value. Trauma patients who **DO NOT** require cervical spine imaging require **ALL** of the following:

- Alert and stable
- **NO** altered level of consciousness
- **NO** distracting injury
- **NO** focal neurological deficit
- **NOT** intoxicated
- **NO** midline spinal tenderness

Website for more information: <https://www.mdcalc.com/calc/703/nexus-criteria-c-spine-imaging#:~:text=The%20NEXUS%20Criteria%20were%20developed,safely%20avoided%20in%20appropriate%20patients.>

Nerve conduction study (NCS) is a test that measures how fast an electrical impulse moves through the nerve and can identify nerve damage.

Neurocutaneous disorders are disorders that affect the brain, spinal cord, organs, skin and bones. The diseases are lifelong conditions that can cause tumors to grow in these areas.

Non-diagnostic is a result that does not lead to a confirmed diagnosis.

Non-union fracture is demonstrated by no healing between two sets of x-rays. Incomplete healing by 6 to 8 months is non-union.

Orthostatic headache is a headache while upright, that is relieved by lying down.

Ossification of the Posterior Longitudinal Ligament (OPLL) is a condition where the ligament, that runs along the back of the bone (vertebral body) and disc, hardens into bone.

Osteomyelitis is an infectious, inflammatory disease of bone. It is often painful, bacterial in origin and may result in the death of bone tissue.

Otorrhea is drainage of liquid from the ear.

Pediatric approximate ages are defined by the US Department of Health (USDH), the Food and Drug Administration (FDA), and the American Academy of Pediatrics (AAP) as the following:

- Infancy, between birth and 2 years of age
- Childhood, from 2 to 12 years of age

- Adolescence, from 12 to 21 years of age, further defined by the AAP into:
 1. Early (ages 11–14 years)
 2. Middle (ages 15–17 years),
 3. Late (ages 18–21 years)
 4. Older ages may be appropriate for children with special healthcare needs.

Radiculopathy is an irritation of or injury to a spinal nerve root (as from being compressed) that typically causes pain, numbness or muscle weakness in the part of the body which is supplied with nerves from that root.

Recurrence is a new occurrence of something that happened or appeared before.

Rheumatoid arthritis (RA) is an autoimmune disease (usually chronic) that is characterized by pain, stiffness, inflammation, swelling and sometimes destruction of the joints.

Rhinorrhea is excessive mucous drainage from the nose.

Sarcoidosis is a chronic disease of unknown cause, that is characterized by the formation of nodules, especially in the lymph nodes, lungs, bones and skin.

Scoliosis is a lateral curvature of the spine of at least 10° with vertebral rotation, presenting as a three-dimensional spinal deformity.

Short segment is a curve in the spinal column that is less than 6 segments.

Spinal cord compression occurs when pressure on the spinal cord prevents the nerves from functioning normally, causing deficits.

Spinal dysraphism is a congenital abnormality that results in an abnormal structure in the spine, including the bony structure, the spinal cord and the nerve roots.

Spondylarthropathy is an inflammatory arthritis affecting the spine.

Spondylosis is a condition in which there is abnormal wear on the cartilage and bones of the neck (cervical vertebrae). It is a common cause of chronic neck pain. Spondylosis is caused by chronic wear on the spine.

Spontaneous intracranial hypotension (SIH) is a condition characterized by cerebrospinal fluid (CSF) hypovolemia due to a noniatrogenic spinal CSF leak, often presenting with orthostatic headache.

Staging in cancer is the process of determining how much cancer is within the body (tumor size) and if it has metastasized (spread).

Stenosis is a narrowing or constriction of the diameter of a bodily passage or orifice.

Surveillance in cancer is the ongoing, timely and systematic collection and analysis of information on new cancer cases, extent of disease, screening tests, treatment, survival and cancer deaths.

Syringomyelia is a chronic progressive disease of the spinal cord associated with sensory disturbances, muscle atrophy and spasticity.

Syrinx is a cerebrospinal fluid-filled cyst which collects inside of the spinal cord or brain stem. A syrinx in the spinal cord is called syringomyelia, and a syrinx in the brain stem is called syringobulbia.

Tethered spinal cord syndrome (TSCS) is a disorder of the nervous system caused by tissue that attaches itself to the spinal cord and limits the movement of the spinal cord.

CT Cervical Spine References

- [1] Agarwal, V., Shah, L.M., . . . Corey, A.S. (2021). ACR Appropriateness Criteria Myelopathy: 2021 Update. *Journal of the American College of Radiology*, 18(5S), S73-S82.
- [2] American College of Radiology. (2024). ACR-ASNR-SPR PRACTICE PARAMETER FOR THE PERFORMANCE OF MYELOGRAPHY AND CISTERNOGRAPHY. *American College of Radiology*. Retrieved: March 2025. <https://gravitas.acr.org/PPTS/GetDocumentView?docId=64+&releaseId=2#>
- [3] American College of Radiology. (2023). ACR Manual on Contrast Media. *American College of Radiology*. Retrieved: May 2024. https://www.acr.org/-/media/ACR/Files/Clinical-Resources/Contrast_Media.pdf
- [4] Basra, M., Patel, H., . . . Posey, A. (2024). Use of Multimodality Imaging in the Evaluation of Patients With Spondyloarthropathies and Sacroiliitis. *Cureus*, 16(3), e57185.
- [5] Beckmann, N.M., West, O.C., . . . Bykowski, J. (2019). ACR Appropriateness Criteria Suspected Spine Trauma. *Journal of the American College of Radiology*, 16(5S), S264-S285.
- [6] Bestic, J.M., Wessell, D.E., . . . Kransdorf, M.J. (2020). ACR Appropriateness Criteria Primary Bone Tumors. *Journal of the American College of Radiology*, 17(5S), S226-S238.
- [7] Booth, T.N., Iyer, R.S., . . . Palasis, S. (2017). ACR Appropriateness Criteria Back Pain—Child. *Journal of the American College of Radiology*, 14(5), S13-S24.
- [8] Czuczman, G.J., Mandell, J.C., . . . Beaman, F.D. (2021). ACR Appropriateness Criteria Inflammatory Back Pain: Known or Suspected Axial Spondyloarthritis: 2021 Update. *Journal of the American College of Radiology*, 18(11S), S340-S360.
- [9] Dobrocky, T., Nicholson, P., . . . Piechowiak, E.I. (2022). Spontaneous intracranial hypotension: searching for the CSF leak. *The Lancet Neurology*, 21(4), 269-380.
- [10] Jones, M.R., Shlobin, N.A. & Dahdaleh, N.S. (2021). Spontaneous Spinal Cerebrospinal Fluid Leak: Review and Management Algorithm. *World Neurosurgery*, 150, 133-139.
- [11] Jones, J.Y., Saigal, G., . . . Karmazyn, B. (2019). ACR Appropriateness Criteria Scoliosis—Child. *Journal of the American College of Radiology*, 16(5), S244-S251.
- [12] Junior, C., Champs, S., . . . Ferrari, A. (2021). Reliability of Evaluation of the Craniocervical Junction by XR, CT and MRI in Patients with Genetic Skeletal Diseases. *Austin Journal of Radiology*, 8(1), 1118.

- [13] Kadom, N., Palasis, S., . . . Karmazyn, B. (2019). ACR Appropriateness Criteria Suspected Spine Trauma-Child. *Journal of the American College of Radiology*, 16(5), S286-S299.
- [14] Khan, M.A., Jennings, J.W., . . . Burns, J. (2023). ACR Appropriateness Criteria Management of Vertebral Compression Fractures: 2022 Update. *Journal of the American College of Radiology*, 20(5), S102-S124.
- [15] McDonald, M.A., Kirsch, C.F.E., . . . Bykowski, J. . (2019). ACR Appropriateness Criteria Cervical Neck Pain or Cervical Radiculopathy. *Journal of the American College of Radiology*, 16(5S), S57-S76.
- [16] Naik, A., Kotecha, H., . . . Issrani, M. . (2021). Canadian C-spine rule (CCR) versus national emergency X- radiography utilization study (NEXUS) for screening cervical spine injury. *International Journal of Orthopaedics Sciences*, 7(1), 787-789.
- [17] Ortiz, A.O., Levitt, A., . . . Corey, A.S. (2021). ACR Appropriateness Criteria Suspected Spine Infection. *Journal of the American College of Radiology*, 18(11S), S488-S501.
- [18] Tam, S.K.P., Chia, J.M., . . . Foroughi, M. (2022). Assessment of patients with a Chiari malformation type I. *Brain and Spine*, 2, 100850.
- [19] Vazirizadeh-Mahabadi, M., Yarahmadi, M. (2023). Canadian C-spine Rule versus NEXUS in Screening of Clinically Important Traumatic Cervical Spine Injuries; a systematic review and meta-analysis. *Archives of Academic Emergency Medicine*, 11(1), e5.
- [20] Watkins, R.G. & Watkins, R.G. (2021). Cervical Disc Herniations, Radiculopathy, and Myelopathy. *Clinics in Sports Medicine*, 40(3), 513-539.

Disclaimer section

Purpose

The purpose of the HealthHelp's clinical guidelines is to assist healthcare professionals in selecting the medical service that may be appropriate and supported by evidence to safely improve outcomes. Medical information is constantly evolving, and HealthHelp reserves the right to review and update these clinical guidelines periodically. HealthHelp reserves the right to include in these guidelines the clinical indications as appropriate for the organization's program objectives. Therefore the guidelines are not a list of all the clinical indications for a stated procedure, and associated Procedure Code Tables may not represent all codes available for that state procedure or that are managed by a specific client-organization.

Clinician Review

These clinical guidelines neither preempt clinical judgment of trained professionals nor advise anyone on how to practice medicine. Healthcare professionals using these clinical guidelines are responsible for all clinical decisions based on their assessment. All Clinical Reviewers are

instructed to apply clinical indications based on individual patient assessment and documentation, within the scope of their clinical license.

Payment

The use of these clinical guidelines does not provide authorization, certification, explanation of benefits, or guarantee of payment; nor do the guidelines substitute for, or constitute, medical advice. Federal and State law, as well as member benefit contract language (including definitions and specific contract provisions/exclusions) take precedence over clinical guidelines and must be considered first when determining eligibility for coverage. All final determinations on coverage and payment are the responsibility of the health plan. Nothing contained within this document can be interpreted to mean otherwise.

Registered Trademarks (®/™) and Copyright (©)

All trademarks, product names, logos, and brand names are the property of their respective owners and are used for purposes of information and/or illustration only. Current Procedural Terminology (CPT)®™ is a registered trademark of the American Medical Association (AMA). No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from HealthHelp.

National and Local Coverage Determination (NCD and LCD)



NOTICE

To ensure appropriate review occurs to the most current NCD and/or LCD, always defer to <https://www.cms.gov/medicare-coverage-database/search.aspx>.

Background

National Coverage Determinations (NCD) and Local Coverage Determinations (LCD) are payment policy documents outlined by the Centers for Medicare and Medicaid Services (CMS) and the government's delegated Medicare Audit Contractors (MACs) that operate regionally in jurisdictions.

CMS introduced variation between different jurisdictions/Medicare Audit Contractors (MACs) and their associated covered code lists with the transition to ICD 10. The variation resulted in jurisdictions independently defining how codes are applied for exclusions, limitations, groupings, ranges, etc. for the medical necessity indications outlined in the NCD and LCD. Due to this variation, there is an inconsistent use/application of codes and coverage determinations across the United States between the different MACs.



A WNS COMPANY

In addition, **WITHOUT** notice, CMS can change the codes that indicate medical necessity and the format of the coverage determinations/associated documents (eg, Articles). This is an additional challenge for organizations to keep up with ongoing, unplanned changes in covered codes and medical necessity indications.

Medical Necessity Codes

Due to the variation in code application between jurisdictions/MACs and that updates can happen without notification, HealthHelp is not able to guarantee full accuracy of the codes listed for any Coverage Determination, and advises that prior to use, the associated Coverage Determination Articles are reviewed to ensure applicability to HealthHelp's programs and any associated NCDs and LCDs.

For Internal Use Only:

11248 11249 11253 11282 11325 11328 11333 11349 11350 11351 11352 11354 11355 11356
11358 11359 11360 11361 11362 11365 11366 11367 11368 11369 11370 11374 11375 11394
11395 11396 11565